

Characterizing Subgroups of Compact Abelian Groups

Dikran Dikranjan* and Kenneth Kunen†

February 1, 2008

Abstract

We prove that every countable subgroup of a compact metrizable abelian group has a characterizing set. As an application, we answer several questions on maximally almost periodic (MAP) groups and give a characterization of the class of (necessarily MAP) abelian topological groups whose Bohr topology has countable pseudocharacter.

1 Introduction

We shall write our abelian groups additively, and we view the circle $\mathbb{T}$ as $\mathbb{R}/\mathbb{Z}$; then, for $z \in \mathbb{T}$, $\|z\|$ is the distance to the nearest integer, so $0 \leq \|z\| \leq 1/2$. All topological groups are assumed to be Hausdorff unless otherwise noted.

For a topological abelian group X , $\widehat{X}$ denotes the group of all continuous characters on X (i.e., the continuous homomorphisms from X to $\mathbb{T}$), and $\widehat{X}$ is given the compact open topology. When X is locally compact, the *Pontryagin Duality Theorem* lets us identify X with $\widehat{\widehat{X}}$; see [19, 22, 27]. For most of the results in this paper, X will be compact, so that $\widehat{X}$ will be discrete.

One can use countable sequences or countable sets of characters to define subgroups of X as follows:

Definition 1.1 *Let X be a topological abelian group:*

- a. [16] *For a sequence $\underline{u} = \langle u_n : n \in \omega \rangle$ of elements of $\widehat{X}$, let $s_{\underline{u}}(X)$ be the set of all $x \in X$ such that $u_n(x) \rightarrow 0$. If $H = s_{\underline{u}}(X)$, we say that $\underline{u}$ characterizes H .*
- b. [20, 21] *For a countably infinite subset B of $\widehat{X}$, let $\mathcal{C}_B(X)$ be the set of all $x \in X$ such that $\langle \varphi(x) : \varphi \in B \rangle$ converges to 0 in $\mathbb{T}$. If $H = \mathcal{C}_B(X)$, we say that B characterizes H .*

*Partially supported by INDAM (GNSAGA)

†Partially supported by NSF Grant DMS-0097881.

Note that $s_{\underline{u}}(X)$ and $\mathcal{C}_B(X)$ are both subgroups of X . We study here which subgroups of X can be characterized as an $s_{\underline{u}}(X)$ or a $\mathcal{C}_B(X)$. Observe that $s_{\underline{u}}(X)$ and $\mathcal{C}_B(X)$ are minor variants of the same notion:

Lemma 1.2 *Let X be any compact topological abelian group.*

1. $\mathcal{C}_B(X) = s_{\underline{u}}(X)$ for any $\underline{u}$ which is a 1-1 enumeration of B .
2. If H is closed and $|X : H|$ is finite, then $H = s_{\underline{u}}(X)$ for some u .
3. Every $\mathcal{C}_B(X)$ is a Haar null set.
4. If $|X : H|$ is infinite and $H = s_{\underline{u}}(X)$, then $H = \mathcal{C}_B(X)$ for some B .

Proof. (1) is clear from the definitions. For (2), $H^\perp = \{\varphi \in \widehat{X} : \varphi(H) = \{0\}\}$ is finite (of size $|X : H|$), and $H = s_{\underline{u}}(X)$ for any $\underline{u} : \omega \rightarrow H^\perp$ which lists each character in $H^\perp$ infinitely often. (3) follows easily from the orthogonality of the characters; see, e.g., [4, 13, 20, 21]. (4) will be proved in Section 2. $\square$

Note also that every $\mathcal{C}_B(X)$ and $s_{\underline{u}}(X)$ is a Borel set (in fact, an $F_{\sigma\delta}$), and every Borel subgroup of finite index is closed and open and of positive Haar measure. By Lemma 1.2, characterization as an $s_{\underline{u}}(X)$ is equivalent to characterization as a $\mathcal{C}_B(X)$ except in the trivial case of subgroups of finite index.

The subgroups $s_{\underline{u}}(X)$ have been studied by many authors, especially in the case $X = \mathbb{T}$, where $\widehat{X}$ is identified with $\mathbb{Z}$ (see [1, 5, 6, 10, 11, 15, 18, 24, 25]). When $x \in \mathbb{T}$ is a non-torsion element, the behavior of the sequences of the form $\langle u_n(x) : n \in \omega \rangle$ is related to Diophantine approximation and dynamical systems (more specifically, the Sturmian flow [26]), as well as to the study of precompact group topologies with converging sequences [4, 7, 16].

Some instances of characterization in $\mathbb{T}$ were described in Armacost's book [1], although this book predates the “characterization” terminology. Specifically, for each prime p , the Prüfer subgroup $\mathbb{Z}_{p^\infty}$ of $\mathbb{T}$ is characterized by $\{p^n : n \in \omega\}$. Also, let $D = \{n! : n \in \omega\}$. Then the set of rational points of $\mathbb{T}$ (i.e., $\mathbb{Q}/\mathbb{Z}$) is properly contained in $\mathcal{C}_D$ (in fact, this $\mathcal{C}_D$ is uncountable). The elements of $\mathcal{C}_D$ are called the *topologically torsion elements* of $\mathbb{T}$ in [1]; they are described further in [14, 15, 17].

It is easy to see that $\mathbb{Q}/\mathbb{Z}$ can be characterized by some set of characters:

Proposition 1.3 *If $B = \{k \cdot n! : 0 < k \leq n \in \omega\}$, then $\mathbb{Q}/\mathbb{Z} = \mathcal{C}_B(\mathbb{T})$.*

It is not always so easy to write down a characterizing set explicitly, but we shall show:

Theorem 1.4 *Every countably infinite subgroup of a compact metrizable abelian group X has a characterizing set.*

This theorem resolves Problem 5.3 from [16]. The proposition and theorem will be proved in Section 2. A number of important special cases of the theorem are already in the literature. In particular, when $X = \mathbb{T}$, it was proved by B  r  , Deshouillers, and S  s [10]. Earlier results by Larcher [25] and Kraaikamp and Liardet [24] relate the characterizability of infinite cyclic subgroups $\langle \alpha \rangle$ of $\mathbb{T}$ to the continued fraction representation of α ; see also [5]. It was proved in [16] that every cyclic subgroup of the group $\widehat{\mathbb{Z}_p^\infty}$ of p -adic integers has a characterizing set. Furthermore, B  r   [9] has proved Theorem 1.4 in the case that the subgroup is dense in X , finitely generated, and torsion-free. Also, Beigl  ock, B  r  , S  s, and Winkler (unpublished) have proved Theorem 1.4 by a different method.

Note that in Theorem 1.4, X must be metrizable (equivalently, second countable), since $\mathcal{C}_B(X)$ always contains $\bigcap_{\varphi \in B} \ker(\varphi)$, and this set will be uncountable if X is not metrizable. We do not know a simple general criterion for deciding whether $H \leq X$ is characterizable. For closed H , this is easy; H is of the form $s_{\underline{u}}(X)$ iff H is a G_δ (see Proposition 1.8 below), and then, applying Lemma 1.2, H is of the form $\mathcal{C}_B(X)$ iff H is a G_δ of infinite index in X . In Section 2, we shall prove the following theorem, which shows that not all F_σ subgroups can be characterized, even in metrizable compact groups:

Theorem 1.5 *Suppose that X is a compact abelian group and $H = \bigcup_n F_n$, where each $F_n \leq X$ is closed and each $F_n \subsetneq F_{n+1}$. Then the following are equivalent:*

- a. $H = \mathcal{C}_B(X)$ for some countably infinite $B \subseteq \widehat{X}$.
- b. $H = s_{\underline{u}}(X)$ for some $\underline{u} : \omega \rightarrow \widehat{X}$.
- c. For some m : X/F_m is metrizable and $|F_{n+1} : F_n|$ is finite for all $n \geq m$.

Next, we give an application of Theorem 1.4 to non-compact groups. If (G, τ) is a topological abelian group, we let (G, τ^+) be its *Bohr modification*. So, τ^+ is the coarsest topology which makes all (continuous) characters of (G, τ) continuous. Clearly, τ^+ is coarser than τ . When τ is clear from context, we refer to (G, τ^+) as G^+ . It is easy to find Hausdorff G for which G^+ is indiscrete, but we are primarily interested in the case where G^+ is also Hausdorff; such G are called *MAP* (*maximally almost periodic*).

A topological space X has *countable pseudocharacter* iff every singleton $\{x\}$ is a G_δ set. This implies that X is a T_1 space, which, in the case of topological groups, is equivalent to being Hausdorff.

Corollary 1.6 *For a MAP topological abelian group G the following are equivalent:*

- a. Every countable $H \leq G$ is of the form $s_{\underline{u}}(G)$ for some $\underline{u} : \omega \rightarrow \widehat{G}$.
- b. Some countable $H \leq G$ is of the form $s_{\underline{u}}(G)$ for some $\underline{u} : \omega \rightarrow \widehat{G}$.
- c. G^+ has countable pseudocharacter.

Proof. (a) $\rightarrow$ (b) is obvious. For (b) $\rightarrow$ (c), note that $K = \bigcap_n \ker(u_n)$ is a G_δ set in G^+ , and $K \leq H$, so K is countable. Since G^+ is Hausdorff, it follows that $\{0\}$ is a G_δ set in G^+ .

For (c) $\rightarrow$ (a): Since $\{0\}$ is a G_δ set in G^+ , there is a countable $\Phi \subseteq \widehat{G}$ such that $\{0\} = \bigcap \{\ker(\varphi) : \varphi \in \Phi\}$. Let $\Delta : G \rightarrow \mathbb{T}^\Phi$ be the evaluation map: $\Delta(x)(\varphi) = \varphi(x)$. Then Δ is 1-1 and continuous. Applying Theorem 1.4 and Lemma 1.2.1, let $\underline{v} = \langle v_n : n \in \omega \rangle$ be a sequence of characters of $\mathbb{T}^\Phi$ such that $\Delta(H) = s_{\underline{v}}(\mathbb{T}^\Phi)$. If $u_n = v_n \circ \Delta$, then $H = s_{\underline{u}}(G)$. $\square$

Note that if one did not assume that G is MAP in Corollary 1.6, either of condition (a) (just using $\{0\} = s_{\underline{u}}(G)$) or condition (c) would imply that G is MAP anyway.

Since having a characterizing sequence (or set) is a rather restrictive property, one can relax it in the following way, following [16]:

Definition 1.7 *For a subgroup H of a topological abelian group X ,*

$$\mathbf{gcl}(H) = \bigcap \left\{ s_{\underline{u}}(X) : \underline{u} : \omega \rightarrow \widehat{X} \text{ \& } H \subseteq s_{\underline{u}}(X) \right\}.$$

H is $\mathbf{g}$ -closed iff $\mathbf{gcl}(H) = H$.

Of course, H is $\mathbf{g}$ -closed whenever $H = s_{\underline{u}}(X)$ for some $\underline{u}$, but a $\mathbf{g}$ -closed subgroup need not be of the form $s_{\underline{u}}(X)$. For example,

Proposition 1.8 *Suppose that H is a closed subgroup of the compact X . Then:*

1. H is $\mathbf{g}$ -closed.
2. $H = s_{\underline{u}}(X)$ for some $\underline{u}$ iff H is a G_δ set.

Proof. For (1), fix $x \notin H$, and then fix $\varphi \in \widehat{X}$ with $\varphi(x) \neq 0$ but $\varphi(H) = \{0\}$. If $u_n = \varphi$ for all n , then $x \notin s_{\underline{u}}(X)$. Thus, $x \notin \mathbf{gcl}(H)$. For (2) $\leftarrow$, if H is a closed G_δ , then $H^\perp = \{\varphi \in \widehat{X} : \varphi(H) = \{0\}\}$ is countable, and then $H = s_{\underline{u}}(X)$ for any $\underline{u} : \omega \rightarrow H^\perp$ which lists each character in $H^\perp$ infinitely often. For (2) $\rightarrow$, let $K = \bigcap_n \ker(u_n)$, and let $\pi : X \rightarrow X/K$ be the natural quotient map. Then X/K is second countable, and $\pi(H)$ is closed and hence a G_δ in X/K , so $H = \pi^{-1}\pi(H)$ is a G_δ in X . $\square$

Likewise, Theorem 1.4 holds only for compact metrizable X , but metrizability is not needed for:

Corollary 1.9 *Every countably infinite subgroup of a compact abelian group X is $\mathbf{g}$ -closed.*

Proof. Fix a countable H and $x \notin H$. Then, let Γ be a countable subgroup of $\widehat{X}$ which separates the elements of $H \cup \{x\}$, let $K = \bigcap \{\ker(\psi) : \psi \in \Gamma\}$, and let $\pi : X \rightarrow X/K$ be the natural quotient map. Then π is 1-1 on $H \cup \{x\}$, and one can apply Theorem 1.4 in X/K to fix $B \subseteq \widehat{X/K}$ with $\pi(H) = \mathcal{C}_B(X/K)$. If $\{v_n : n \in \omega\}$ is a 1-1 listing of B and $u_n = v_n \circ \pi$, then $H \leq s_{\underline{u}}(X)$ and $x \notin s_{\underline{u}}(X)$. $\square$

This generalizes Theorem 1.2 of [16], which proved the result for cyclic subgroups, and it answers Problem 5.1 and Question 5.2 of [16]. Note that Corollary 1.9 actually holds for all MAP X , since one can apply the corollary in the Bohr compactification, $b(X)$. We do not know if every F_σ subgroup of a compact X is $\mathfrak{g}$ -closed, although, in contrast to Theorem 1.5,

Proposition 1.10 *Suppose that X is a compact abelian group and $H = \bigcup_n F_n$, where each $F_n \leq X$ is closed and each $F_n \leq F_{n+1}$. Then H is $\mathfrak{g}$ -closed.*

Proof. Fix $x \in X \setminus H$. Since F_n is closed, we can choose $u_n \in \widehat{X}$ such that $F_n \leq \ker(u_n)$ and $\|u_n(x)\| \geq 1/4$. Then $H \leq s_{\underline{u}}(X)$ and $x \notin s_{\underline{u}}(X)$. $\square$

$F_{\sigma\delta}$ subgroups certainly need not be $\mathfrak{g}$ -closed. In fact, whenever X is infinite, there is an $F_{\sigma\delta}$ subgroup H such that H is a Haar null set and $\mathfrak{gcl}(H) = X$; see [21], which extends earlier arguments in [4, 20].

The next corollary says that every totally bounded countable metrizable group topology can be “encoded” by means of a *single* τ -convergent zero-sequence:

Corollary 1.11 *Let (G, τ) be a countable totally bounded metrizable abelian group. Then there exists a sequence $\underline{u} = \langle u_n : n \in \omega \rangle$ with $u_n \rightarrow 0$ in (G, τ) such that τ is the finest totally bounded group topology on G in which $u_n \rightarrow 0$.*

Proof. Let G_d denote the group G with the discrete topology, and let $X = \widehat{(G_d)}$. Let $H \leq X$ be the set of $\varphi : G \rightarrow \mathbb{T}$ which are τ -continuous. Then H is countable and X is metrizable. Identifying G_d with $\widehat{X}$, apply Theorem 1.4 to fix $u_n \in G$ with $H = s_{\underline{u}}(X)$. Now, suppose that τ' is a strictly finer totally bounded group topology on G . Then there is a τ' -continuous character x of G which is not τ -continuous, so that $x \in X$ but $x \notin H$. Then $x(u_n) = u_n(x) \not\rightarrow 0$ in $\mathbb{T}$, so $u_n \not\rightarrow 0$ in (G, τ') . $\square$

It is easy to see that countability is necessary; that is, if (G, τ) is a totally bounded metrizable abelian group such that τ is the finest totally bounded group topology on G in which some sequence $u_n \rightarrow 0$, then G is countable.

Corollary 1.11 should be compared to the fact that the finest *Hausdorff* group topology that makes a given sequence converge to 0 is never Fréchet-Urysohn [28]. In the case $G = \mathbb{Z}$, the encoding sequence $\underline{u}$ can often be displayed explicitly. For example, if τ is the p -adic topology, one can take $u_n = p^n$. If τ is the profinite topology, then one can let $\underline{u}$ enumerate $\{k \cdot n! : 0 \leq k \leq n \in \omega\}$ by Proposition 1.3.

2 The Proofs

We begin by relating characterizable subgroups of N to characterizable subgroups of X , where N is a subgroup of X :

Lemma 2.1 *Suppose that $H \leq N \leq X$, where X is a compact abelian group, N is closed in X , X/N is metrizable, and $H = \mathcal{C}_B(N)$ for some countably infinite $B \subseteq \widehat{N}$. Then $H = \mathcal{C}_D(X)$ for some countably infinite $D \subseteq \widehat{X}$.*

Proof. Let $N^\perp = \{\varphi \in \widehat{X} : \varphi(N) = \{0\}\}$, which is countable.

If $|N^\perp| = \aleph_0$: For $\varphi \in B$, choose $\tilde{\varphi} \in \widehat{X}$ such that $\tilde{\varphi}|N = \varphi$. Let $D = \{\tilde{\varphi} : \varphi \in B\} \cup N^\perp$. Then $\mathcal{C}_D^X \cap N = \mathcal{C}_B^N$, and if $y \in X \setminus N$, then $y \notin \mathcal{C}_{N^\perp}^X$, so $y \notin \mathcal{C}_D^X$.

If $|N^\perp| = k < \aleph_0$: Then $|X : N| = k$. Let $D = \{\psi \in \widehat{X} : \psi|N \in B\}$. For $\varphi \in B$, $|\{\psi \in D : \psi|N = \varphi\}| = k$, so $\mathcal{C}_D^X \cap N = \mathcal{C}_B^N$. Now, fix $y \in X \setminus N$, and fix any $\alpha \in N^\perp$ such that $\alpha(y) \neq 0$. Note that $\psi \in D$ iff $\alpha + \psi \in D$, so $\langle \psi(y) : \psi \in D \rangle$ cannot converge to 0. $\square$

Proof of Lemma 1.2.4. We have X compact, $H \leq X$, $|X : H|$ infinite, and $H = s_{\underline{u}}(X)$. We shall produce a countably infinite $D \subseteq \widehat{X}$ such that $H = \mathcal{C}_D(X)$. Let $N = \overline{H}$ and let $K = \bigcap \{\ker(u_n) : n \in \omega\}$. Then $K \leq H \leq N \leq X$, and X/K is metrizable. If H is closed in X , then X/H is metrizable and infinite, so $H = \mathcal{C}_{H^\perp}(X)$.

Now, assume that H is not closed, so $H \neq N$. Let $v_n = u_n|N$, so $H = s_{\underline{v}}(N)$. Let $B = \{v_n : n \in \omega\} \subseteq \widehat{N}$. Then B is infinite (since otherwise H would be closed), and $\{n : u_n = \psi\}$ is finite for each non-zero $\psi \in \widehat{N}$ (since $N = \overline{H}$), so $H = \mathcal{C}_B(N)$. We are now done by Lemma 2.1. $\square$

Proof of Proposition 1.3. It is clear that $\mathbb{Q}/\mathbb{Z} \subseteq \mathcal{C}_B(\mathbb{T})$. Now, any $x \in [0, 1]$ can be written in a factorial expansion as $x = \sum_{n=1}^{\infty} c_n/(n+1)!$, where the integers c_n satisfy $0 \leq c_n \leq n$. Note that $\sum_{n=k}^{\infty} n/(n+1)! = 1/k!$. Thus, if $x \notin \mathbb{Q}$, then $c_n \notin \{0, n\}$ for infinitely many n . For any n , we have $n!x \equiv y_n \pmod{1}$, where $y_n = n! \sum_{m=n}^{\infty} c_m/(m+1)!$. When $c_n \notin \{0, n\}$,

$$\frac{1}{n+1} = n! \left[\frac{1}{(n+1)!} \right] \leq y_n \leq n! \left[\frac{n-1}{(n+1)!} + \frac{1}{(n+1)!} \right] = \frac{n}{n+1} ,$$

so we can find a k with $0 < k \leq n$ such that $\|kn!x\| = \|kn!y_n\| \geq 1/4$. When $x \notin \mathbb{Q}$, there are infinitely many such k, n , so $x \notin \mathcal{C}_B(\mathbb{T})$. $\square$

We now prove three lemmas for Theorem 1.4. As is common in metric spaces, $B(x; \varepsilon)$ denotes an open ball, and $N(F; \varepsilon)$ denotes the open set $\bigcup \{B(x; \varepsilon) : x \in F\}$.

Lemma 2.2 *Let $(X; d)$ be a metric space, F any countably infinite subset of X , and F_n finite subsets such that $F_n \nearrow F$. Then there are positive $\varepsilon_n \searrow 0$ such that for all $x \notin F$, there are infinitely many n with $x \notin N(F_n; \varepsilon_n)$.*

Proof. Choose $\varepsilon_n \searrow 0$ so that $2\varepsilon_n < d(u, v)$ whenever u, v are distinct elements of F_{n+1} . Now, fix m and assume that $x \in N(F_n; \varepsilon_n)$ for all $n \geq m$; we show that $x \in F$. For each $n \geq m$, choose $u_n \in F_n$ such that $d(x, u_n) < \varepsilon_n$. Then $d(u_n, u_{n+1}) \leq d(x, u_n) + d(x, u_{n+1}) < \varepsilon_n + \varepsilon_{n+1} \leq 2\varepsilon_n$, so $u_n = u_{n+1}$. Thus, the u_n are all equal to some $u \in F$, and $x = u$. $\square$

If G is a discrete group, then G is dense in its Bohr compactification; if this fact is stated in terms of the compact group $X = \widehat{G}$, we get the following lemma, which generalizes an old result of Kronecker; see [12, p. 1188], [22, Cor. 26.16], or [23].

Lemma 2.3 *Let X be a compact abelian group, $x_1, \dots, x_t$ a finite list of elements of X , $\psi : X \rightarrow \mathbb{T}$ a possibly discontinuous homomorphism, and $\delta > 0$. Then there is a $\varphi \in \widehat{X}$ such that $\|\psi(x_i) - \varphi(x_i)\| < \delta$ for $i = 1, \dots, t$.*

Using this, one can give a direct proof of Corollary 1.9, without using Theorem 1.4. We have $H < X$, where H is countable and X is compact. Fix $x \notin H$ and list H as $\{e_n : n \in \omega\}$. For each n , $x \notin \langle e_0, e_1, \dots, e_n \rangle$ implies that there is a homomorphism $\psi_n : X \rightarrow \mathbb{T}$ such that $\|\psi_n(x)\| > 1/4$ and $\ker(\psi_n)$ contains $e_0, e_1, \dots, e_n$. By Lemma 2.3, there is a $u_n \in \widehat{X}$ such that $\|u_n(x)\| > 1/4$ and $\|u_n(e_j)\| < 1/n$ for $j = 0, 1, \dots, n$. Then $H \leq s_{\underline{u}}(X)$ and $x \notin s_{\underline{u}}(X)$.

Definition 2.4 *Let X be any topological abelian group and $E \subseteq X$. The m -quasi-convex hull $\text{qc}_m(E)$ of E is the set*

$$\left\{ x \in X : \forall \varphi \in \widehat{X} \left[\forall e \in E \left[\|\varphi(e)\| \leq 2^{-m-2} \right] \implies \|\varphi(x)\| \leq 1/4 \right] \right\}$$

$$\text{qc}(E) = \text{qc}_0(E).$$

The *quasi-convex hull* $\text{qc}(E)$ is discussed in [2, 3]. Note that $\text{qc}_m(E)$ gets bigger as m gets bigger, and $E \subseteq \text{qc}_0(E)$.

Lemma 2.5 *Let $m \in \omega$ and let E be a finite subset of a compact abelian group X . Then:*

- a. $\text{qc}_m(E) \subseteq \langle E \rangle$.
- b. $\text{qc}_m(E)$ is finite.

Proof. For (a), fix $x \notin \langle E \rangle$. There exists a possibly discontinuous homomorphism ψ such that $\psi(E) = \{0\}$ and $\|\psi(x)\| > 1/4$. Apply Lemma 2.3 to find a $\varphi \in \widehat{X}$ with $|\varphi(e)| < 2^{-m-2}$ for each $e \in E$ and $\|\varphi(x)\| > 1/4$, so that $x \notin \text{qc}_m(E)$.

For (b), let $H = \langle E \rangle$. Then H is an internal direct product of cyclic groups; say $H \cong \langle a_1 \rangle \times \cdots \times \langle a_n \rangle$, where $a_1, \dots, a_n \in H$. For each j , the order $o(a_j)$ is either a positive integer or ∞ . Fix a positive integer $M \geq \max\{o(a_j) : o(a_j) \neq \infty\}$ such that also each $e \in E$ is of the form $\sum_{j=1}^n \mu_j a_j$, where each $|\mu_j| \leq M$. Let F_0 be the set of all elements $\sum_{j=1}^n \nu_j a_j$ such that each $|\nu_j| \leq 2^{m+1}M$. Then F_0 is finite, and we show $\text{qc}_m(E) \subseteq F_0$.

Fix $x = \sum_{j=1}^n \nu_j a_j \notin F_0$, with each $|\nu_j| < o(a_j)$. Then, fix ℓ such that $|\nu_\ell| > 2^{m+1}M$. Then $o(a_\ell) > |\nu_\ell| > M$, so $o(a_\ell) = \infty$. We can then define $\psi \in \text{Hom}(H, \mathbb{T})$ so that $\psi(\sum_{j=1}^n \mu_j a_j) = \mu_\ell / (2\nu_\ell) \pmod{1}$ for all $\sum_{j=1}^n \mu_j a_j \in H$. Then $\psi(x) = 1/2$, but for each $e = \sum_{j=1}^n \mu_j a_j \in E$, we have $\|\psi(e)\| < M \cdot 1/(2^{m+2}M) = 2^{-m-2}$. By Lemma 2.3, there is a $\varphi \in \widehat{X}$ such that $\|\varphi(x)\| > 1/4$, but $\|\varphi(e)\| < 2^{-m-2}$ for each $e \in E$. Thus, $x \notin \text{qc}_m(E)$. $\square$

This lemma actually holds for all MAP X , since one can apply the lemma in $\mathbf{b}(X)$. For $m = 0$, this lemma was proved in [2] (see Lemma 7.10 and Theorem 7.11).

The inclusion $\text{qc}_m(E) \subseteq \langle E \rangle$ can fail when E is infinite. For example, $\text{qc}_m(E) = \overline{E}$ for every $m \in \omega$ whenever $E = \langle E \rangle \leq X$.

Proof of Theorem 1.4. Let X be a compact metric abelian group with $G = \widehat{X}$. Let E a countably infinite subgroup of X . In view of Lemma 2.1, we may assume that E is dense in X . Let $E_n \nearrow E$, with each E_n finite. Let $F_n = \text{qc}_n(E_n)$. Then $E_n \subseteq F_n \subseteq E$, $F_n \nearrow E$, and F_n is finite by Lemma 2.5. Let d be a metric for X . Applying Lemma 2.2, choose positive $\varepsilon_n \searrow 0$ such that for all $x \notin E$, there are infinitely many n with $x \notin N(F_n; \varepsilon_n)$.

Let $A_n = \{\varphi \in \widehat{X} : \forall e \in E_n [\|\varphi(e)\| \leq 2^{-n-2}]\}$. Applying the definition of $\text{qc}_n(E_n)$,

$$\forall x \notin F_n \exists \varphi \in A_n [\|\varphi(x)\| > 1/4] \quad .$$

Hence, $\bigcup_{\varphi \in A_n} \{x \in X : \|\varphi(x)\| > 1/4\} \supseteq X \setminus N(F_n; \varepsilon_n)$. Since $X \setminus N(F_n; \varepsilon_n)$ is compact, we can choose a finite $B_n \subseteq A_n$ such that

$$\forall x \notin N(F_n, \varepsilon_n) \exists \varphi \in B_n [\|\varphi(x)\| > 1/4] \quad .$$

Set $B = \bigcup_{n \in \omega} B_n$ and note that B must be infinite, since (using $\overline{E} = X$)

$$\forall \varphi \neq 0 [\#\{n : \varphi \in A_n\} < \aleph_0] \quad . \quad (*)$$

Clearly, $E \subseteq \mathcal{C}_B$. To prove that $E = \mathcal{C}_B$ fix $x \notin E$. Then $x \notin N(F_n; \varepsilon_n)$ for infinitely many n . For these n , choose $\varphi_n \in B_n$ so that $\|\varphi_n(x)\| > 1/4$. By (*), $\{\varphi_n : x \notin N(F_n; \varepsilon_n)\}$ is infinite, so $x \notin \mathcal{C}_B$. $\square$

Next, we prove three lemmas for Theorem 1.5.

Lemma 2.6 *Suppose that X is a compact abelian group, F a closed subgroup, and B a countably infinite subset of $\widehat{X}$. Then $F \leq \mathcal{C}_B$ iff $F \leq \ker(\varphi)$ for all but finitely many $\varphi \in B$.*

Proof. For the non-trivial direction, assume that $F \leq \mathcal{C}_B$ but $F \not\leq \ker(\varphi)$ for infinitely many $\varphi \in B$. Since $\mathcal{C}_B$ gets bigger as B gets smaller, we may assume that $\varphi|F \neq 0$ for all $\varphi \in B$, and that one of the following two cases holds:

Case I: The $\varphi|F = \psi \in \widehat{F}$ for all $\varphi \in B$: But then any $x \in F \setminus \ker(\psi)$ will be in $F \setminus \mathcal{C}_B$, contradicting $F \leq \mathcal{C}_B$.

Case II: The $\varphi|F$ for $\varphi \in B$ are all different. Let $D = \{\varphi|F : \varphi \in B\}$. Then $\mathcal{C}_B \cap F = \mathcal{C}_D$, which is a null set in F by Lemma 1.2, contradicting $F \leq \mathcal{C}_B$. $\square$

Lemma 2.7 *Suppose that X is a compact metric abelian group, with an invariant metric d . Let $H \leq X$, where $H = \bigcup_n F_n$, each F_n is closed, and each $F_n < F_{n+1} < X$. Assume that $y_n \in F_{n+1} \setminus F_n$ are chosen so that each $d(y_{n+1}, 0) \leq (d(y_n, F_n))/3$. Define $x = \sum_{k=0}^{\infty} y_k$. Then $x \notin H$.*

Proof. By induction on k , we have $d(y_{n+k}, 0) \leq (d(y_n, F_n)) \cdot 3^{-k}$ for all $n \geq 0$ and $k > 0$. This shows that the sum defining x really converges, and also lets us define $x_n = \sum_{k=n}^{\infty} y_k$; so $x = x_0$. Now, fix n , and we show that $x \notin F_n$. Since $(x - x_n) \in F_n$, it is sufficient to show that $x_n \notin F_n$. Now $y_n \notin F_n$ and $x_n = y_n + x_{n+1}$. Also, $d(x_{n+1}, 0) \leq \sum_{k=1}^{\infty} d(y_{n+k}, 0) \leq d(y_n, F_n) \cdot \sum_{k=1}^{\infty} 3^{-k} = d(y_n, F_n)/2$. Thus, $d(x_n, F_n) \geq d(y_n, F_n) - d(x_{n+1}, 0) \geq d(y_n, F_n)/2 > 0$. $\square$

Finally, we need the (trivial) converse to Lemma 2.1:

Lemma 2.8 *Suppose that $H \leq N \leq X$, where X is a compact abelian group, N is closed in X , H is not closed in X , and $H = \mathcal{C}_D(X)$ for some countably infinite $D \subseteq \widehat{X}$. Then $H = \mathcal{C}_B(N)$ for some countably infinite $B \subseteq \widehat{N}$.*

Proof. By Lemma 1.2.1, $H = s_{\underline{u}}(X)$, where each $u_n \in \widehat{X}$. But then also $H = s_{\underline{v}}(N)$, where each $v_n = u_n|N \in \widehat{N}$. We now get B by Lemma 1.2.4. $\square$

Proof of Theorem 1.5. (a) $\leftrightarrow$ (b) follows from Lemma 1.2, since $|X : H|$ is infinite. For (c) $\rightarrow$ (a), we may, replacing X by X/F_m assume that $F_m = \{0\}$. But then H is countable, so the result follows by Theorem 1.4.

For (a) $\rightarrow$ (c): Assume (a), and let $K = \bigcap \{\ker(\varphi) : \varphi \in B\}$. Then $K \leq H$ and X/K is metrizable. Observe that $K \leq F_m$ for some m ; otherwise, we have $K \cap F_n \not\leq K$, where each $K \cap F_n \not\leq K$. If some $|K : K \cap F_n|$ is finite, this is clearly a contradiction, while if all $|K : K \cap F_n|$ are infinite, this is a contradiction by the Baire Category Theorem.

We are now done if we can show that $|F_{n+1} : F_n|$ is finite for all but finitely many n ; so, assume that this is false and we shall derive a contradiction. Re-indexing, we

may assume that $|F_{n+1} : F_n|$ infinite for each n , and we may assume that $F_0 = \{0\}$. We may also assume that $X = \overline{H}$; if not, replace X by $N = \overline{H}$ and apply Lemma 2.8. Then $\widehat{X} = F_0^\perp > F_1^\perp > F_2^\perp \cdots$, and $\bigcap_n F_n^\perp = \{0\}$.

We may assume that X is metrizable; otherwise replace X by X/K . We may assume that $0 \notin B$, so that $B = \bigcup_n B_n$, where $B_n = B \cap F_n^\perp \setminus F_{n+1}^\perp$. By Lemma 2.6, each B_n is finite.

We shall now produce an $x \in \mathcal{C}_B(X) \setminus H$, contradicting $H = \mathcal{C}_B(X)$.

Let d be an invariant metric for X . Inductively choose $y_n \in F_{n+1} \setminus F_n$ so that $\|\varphi(y_n)\| \leq 2^{-n}$ for all $\varphi \in B_0 \cup \cdots \cup B_n$ and each $d(y_{n+1}, 0) \leq (d(y_n, F_n))/3$; this is possible because each F_n is nowhere dense in F_{n+1} . Let $x = \sum_{k=0}^\infty y_k$. Then $x \notin H$ by Lemma 2.7.

If $\varphi \in B_n$, then $\|\varphi(x)\| \leq \sum_{k=0}^\infty \|\varphi(y_k)\| = \sum_{k=n}^\infty \|\varphi(y_k)\| \leq \sum_{k=n}^\infty 2^{-k} = 2^{1-n}$. Thus $\langle \varphi(x) : \varphi \in B \rangle$ converges to 0 in $\mathbb{T}$. $\square$

References

- [1] D. L. Armacost, *The structure of locally compact Abelian groups*, Monographs and Textbooks in Pure and Applied Mathematics, **68**, Marcel Dekker, Inc., New York, 1981.
- [2] L. Aussenhofer, *Contributions to the duality theory of abelian topological groups and to the theory of nuclear groups*, Dissertationes Math. (Rozprawy Mat.) 384 (1999).
- [3] W. Banaszczyk, *Additive subgroups of topological vector spaces*, Lecture Notes in Mathematics, 1466. Springer-Verlag, Berlin, 1991.
- [4] G. Barbieri, D. Dikranjan, C. Milan, H. Weber, *Answer to Raczkowski's questions on convergent sequences*, Top. Appl. **132** (2003), 89–101.
- [5] G. Barbieri, D. Dikranjan, C. Milan and H. Weber, *Topological torsion related to some recursive sequences of integers*, submitted.
- [6] G. Barbieri, D. Dikranjan, C. Milan and H. Weber, *$\mathfrak{t}$ -dense subgroups of topological Abelian groups*, submitted.
- [7] G. Barbieri, D. Dikranjan, C. Milan and H. Weber, *Convergent sequences in precompact group topologies*, submitted.
- [8] A. B  r  , *Characterizing sets for subgroups of compact groups I: a special case*, preprint.
- [9] A. B  r  , *Characterizing sets for subgroups of compact groups II: general case*, preprint.

- [10] A. B  r  , J.-M. Deshouillers and V. S  s, *Good approximation and characterization of subgroups of $\mathbb{R}/\mathbb{Z}$* , Studia Sci. Math. Hungar. **38** (2001), 97–113.
- [11] A. B  r   and V. S  s, *Strong characterizing sequences in simultaneous Diophantine approximation*, J. Number Theory **99** (2003), 405–414.
- [12] W. Comfort *Topological groups*, in: Handbook of Set-Theoretic Topology, edited by K. Kunen and J. E. Vaughan, North Holland, Amsterdam · New York · Oxford (1984), p.1184.
- [13] W.W. Comfort, F.J. Trigos-Arrieta and T.S. Wu, *The Bohr compactification, modulo a metrizable subgroup*, Fundamenta Math. **143** (1993), 119–136, Correction: **152** (1997), 97–98.
- [14] D. Dikranjan, *Topologically torsion elements of topological groups*, Topology Proc., **26**, 2001–2002, pp. 505–532.
- [15] D. Dikranjan and R. Di Santo, *On Armacost’s quest on topologically torsion elements*, Communications Algebra, **32** (2004), 133–146.
- [16] D. Dikranjan, C. Milan and A. Tonolo *A characterization of the MAP abelian groups*, J. Pure Appl. Algebra (2004) to appear.
- [17] D. Dikranjan, Iv. Prodanov and L. Stoyanov, *Topological groups. Characters, dualities and minimal group topologies*, Monographs and Textbooks in Pure and Applied Mathematics **130**, Marcel Dekker, Inc., New York, 1990.
- [18] H.G. Eggleston, *Sets of fractional dimensions which occur in some problems of number theory*, Proc. London Math. Soc. (2) **54**, (1952) 42–93.
- [19] G. B. Folland, *A Course in Abstract Harmonic Analysis*, CRC Press, 1995.
- [20] J. Hart and K. Kunen, *Limits in function spaces and compact groups*, Topology Appl., to appear.
- [21] J. Hart and K. Kunen, *Limits in compact abelian groups*, to appear.
- [22] E. Hewitt and K. Ross, *Abstract Harmonic Analysis*, Volume I, Springer-Verlag, Berlin-G  ttingen-Heidelberg 1979.
- [23] E. Hewitt and H. Zuckerman, *A group-theoretic method in approximation theory*, Ann. of Math. (2) **52** (1950), 557–567.
- [24] C. Kraaikamp and P. Liardet, *Good approximations and continued fractions*, Proc. Amer. Math. Soc. **112** (1991), 303–309.

- [25] G. Larcher, *A convergence problem connected with the continued fractions*, Proc. Amer. Math. Soc., **103** (1988), 718–722.
- [26] K. Petersen and L. Shapiro, *Induced flows*, Trans. Amer. Math. Soc. **177** (1973), 375–390.
- [27] W. Rudin, *Fourier Analysis on Groups*, Interscience Publishers, 1962.
- [28] E. G. Zelenyuk and I. V. Protasov, *Topologies on abelian groups*, Math. USSR Izvestiya **37** (1991), 445–460. Russian original: Izvestia Akad. Nauk SSSR **54** (1990), 1090–1107.

DIPARTIMENTO DI MATEMATICA E INFORMATICA, UNIVERSITÀ DI UDINE,
VIA DELLA SCIENZE 206, 33100 UDINE, ITALY
Email address: dikranja@dimi.uniud.it

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON,
WI 53706, USA
Email address: kunen@math.wisc.edu
URL: <http://www.math.wisc.edu/~kunen>